

Toward a Fire and Haze Early Warning System for Southeast Asia

Jin Ho YOO^a, Jaepil CHO^a✉, Saji HAMEED^b, Robert FIELD^c, Kok Foo KWAN^d, Israr ALBAR^e

APN Project Reference: ARCP2013-10CMY-Yoo / Received: January 2016 / Published online: March 2016

Available online at APN E-Lib: <http://www.apn-gcr.org/resources>

HIGHLIGHTS

- Four different downscaling methods were developed and integrated into the prototype of EWS in order to improve the predictability.
- Long-term predictability of monthly precipitation for the four regions within Borneo Island was evaluated.
- APCC led a two-day workshop in Malaysia, including hands-on training sessions on statistical downscaling and prototype.
- Needs assessment for early warning information was conducted through field surveys with resource managers.
- Monthly precipitation forecasts for dry season (August to October) over 4 provinces in Borneo Island showed good predictability less than four-month lead time by showing temporal correlation coefficients (TCCs) greater than 0.5 in all provinces.

ABSTRACT Smoke haze from forest fires is among Southeast Asia's most serious environmental problems and there is a clear need for a fire and haze early warning system (EWS) for the region. APEC Climate Center (APCC) has been collecting monthly dynamic prediction data produced by 16 institutions and has been producing 6-month lead multi-model ensemble (MME) climate forecasts every month. In this study, we developed four different statistical downscaling methods and assessed the forecast skill of the integrated forecast system over four provinces in Borneo Island. We developed a EWS prototype in which three-month precipitation (August to October) is predicted during April to July and the forecasted precipitation amount is then translated into four fire danger ratings based on the relationship between precipitation amount and CO₂ emission. A needs assessment for early warning information was conducted through field surveys with resource managers at three provinces in Indonesia. A two-day workshop was held for the improvement of the EWS. Finally, the forest fire early warning information on Borneo Island created using the EWS will be provided through the hosting server in APCC.

KEYWORDS *fire danger; seasonal forecasts; statistical downscaling; dynamical downscaling; seasonal drought*

1. Introduction

Smoke haze from forest fires is among Southeast Asia's most serious environmental problems. Severe burning in Indonesia occurs only during years with anomalously low rainfall. Monitoring for these conditions is important, but has limited effectiveness because the burning is opportunistic. As a result, measures to prevent these fires and mitigate their impacts remains limited by the absence of long-lead early warning system (EWS). Severe burning conditions, therefore, need to be forecast weeks to months in advance for any prevention to be effective. In this context, little of the progress made in seasonal forecasting has been applied to fire early warning in Indonesia and there is a clear need for a fire and haze EWS for the region. The project builds upon current fire danger rating systems by providing forecasts at a longer lead-time using seasonal forecast data maintained at APCC, a time-scale that is more relevant and useable for decision makers. The final objective of the project is to develop a prototype of fire danger EWS by considering field survey results and conducting a training workshop.

a APEC Climate Center, 12 Centum 7-ro, Haeundae-gu, Busan, 612-020, Republic of Korea

b Aizu University, street, Aizu-Wakamatsu City, Fukushima-ken 965-8580, Japan

c Columbia University, 116th Street and Broadway, New York, 10027, USA

d Malaysian Meteorological Department, Petaling Jaya, Selangor, 47301, Malaysia

e Department of Forestry, Gedung Manggala Wanabakti Blok, Jakarta, 10207, Indonesia

✉ Corresponding author. Email: jpcho89@gmail.com;

Tel: +82-51-745-3994, Fax: +82-51-745-3999.

2. Methodology

An EWS for forest fire was developed based on an open source license for further training workshops and free distribution of the developed prototype. The overall procedures for development of EWS prototype include 1) construct statistical downscaling model for forecasting monthly area-average precipitation amount for each region, 2) determine number of categories and corresponding ranges of fire danger rating system based on the relationship between total three-month precipitation amount and CO₂ emission, and 3) forecast probabilistic fire danger ratings based on predicted precipitation (Figure 1).

Regarding the statistical downscaling, four different downscaling methods in accordance with the degree of utilising the seasonal climate prediction information were selected for developing the EWS. These methods are: the Simple Bias Correction (SBC), the Moving Window Regression (MWR), the Climate Index Regression (CIR), and the Integrated Time Regression (ITR). SBC is a forecast-based direct downscaling method, which uses GCM's prediction data after adjusting the monthly mean of predicted data. For example, if the precipitation prediction data on a specific region is needed, SBC directly uses the grid values of precipitation variables, which are produced from GCMs over the given area. The systematic bias is adjusted for making the monthly average of prediction the same as the average of observation for the same period. Table 1 shows the selected dynamical prediction models used in the study. If there are limitations in directly predicting target variables such

as precipitation in the target area, the MWR method uses the oceanic and atmospheric circulation variables as predictors to improve the seasonal prediction predictability in the target area (eg. Kang, Park, Hameed, & Ashok, 2009; Kang, Hur, & Ahn, 2014). As a result, MWR is a forecast-based indirect statistical downscaling method, which uses the simultaneous proxy variables produced by GCMs as predictors of regression model when high correlation exists between proxy variables and regional target variables. CIR is an observation-based indirect statistical downscaling method that can be used when there is a high correlation between global climate indices and regional target variables with lag time (eg. Kim, Kim, & Lee, 2007; Kim & Kim, 2010). Twenty five climate indices which are updated monthly from NOAA (<http://www.esrl.noaa.gov/psd/data/climateindices/list/>) and APCC (<http://www.apcc21.org/ser/indic.do?lang=en>) were used for real time operation of CIR method. In this case, lag time between the monthly precipitation and indices should be larger than the lead-time. The CIR method is similar to the MWR method in that both methods indirectly utilise the correlation between regional target variables and global scale climate variables related to oceanic and atmospheric circulation. There is a difference between the CIR and MWR methods when selecting predictors to forecast future seasonal target variable values. While the MWR method uses the forecasted climate information, the CIR method uses the observed information from a few months before taking into account the lag time. ITR is an indirect statistical downscaling method that uses both forecast and observation-based predictors from the




FIGURE 1. Schematic diagram of Early Warning System (EWS) prototype.

Model	Institution	Raw Resolution	Ensemble Size
CANCM3	Meteorological Service of Canada (Canada)	T63L31 (AGCM) 1.41° × 0.94° L40 (OGCM)	10
CANCM4	Meteorological Service of Canada (Canada)	75.37	10
NASA	National Aeronautics and Space Administration (USA)	220.1	10
NCEP	Climate Prediction Center - NCEP/ NWS/NOAA (USA)	11.41	17
PNU	Pusan National University (Republic of Korea)	320.41	4
POAMA	Centre for Australian Weather and Climate Research/ Bureau of Meteorology (Australia)	T47L17 (AGCM) 0.5–1.5°LAT × 2°LON, L25 (OGCM)	30

TABLE 1. Description of dynamical seasonal prediction models used in the study.

MWR and CIR methods, respectively. As a result, it can be used only when the MWR and CIR methods simultaneously select predictors for a particular target period.

We used the same regions from previous research by Field and Shen (2008) for developing and evaluating the statistical downscaling methods. The regions include Southern Sumatra (SSUM), Central Sumatra (CSUM), Eastern Kalimantan (EKAL) and Southern Kalimantan (SKAL). However, we decided the administrative boundary for managerial purpose of the EWS and four provinces in Borneo Island were used in this study (Figure 2).

Second, an analysis of the threshold levels for the study regions was conducted in order to translate the predicted precipitation amount to the fire danger ratings. If the amount of precipitation dips below the threshold level, this predicts an increased risk for severe burning, carbon emissions, and transboundary haze. It is necessary to connect the forecasted precipitation to the possible EWS index based on region-specific threshold level. We used the relationship between region-average ASO precipitation and carbon emission data. The region-average monthly precipitation and carbon emission data were derived from APHRODITE's Water Resources ([http://](http://www.chikyu.ac.jp/precip/)

www.chikyu.ac.jp/precip/) and Global Fire Emissions Database (<http://www.globalfiredata.org/>) webpages, respectively. At first, we attempted to determine the ranges for each category using a segmented regression method. However, the resulting threshold precipitation was too low, which increased the likelihood of extreme carbon emissions being predicted due to scattered data. As a result, we set the threshold value manually based on the time series of three-month accumulated monthly precipitation and carbon emissions.

Based on an earlier version of the prototype, APCC led a workshop including hands-on training sessions on statistical downscaling and the prototype. After the training, we improved the predictor selection algorithm for the MWR and CIR methods in order to avoid overfitting in real-time forecasts. The concepts of both cross-validation and split-validation were applied in order to prevent overfitting problems. The Leave-one-out Cross-Validation (LOOCV) technique was applied to the observation period (1983–2013). In other words, when predicting target variables for a specific target period (year/month), all predictors for the same target period are removed from the model construction procedure in order to reproduce the same conditions as real time forecasting. For example, when predicting




FIGURE 2. Selected regions for developing statistical downscaling methods and early warning system.

for January 1983, only predictors from January 1984 to 2013 are utilised in constructing the regression model. Predictions are made in the same way for the rest of the simulation period. For each cross-validation process, the split validation approach was applied, and then the best predictors that showed consistent performance for both training and verification periods were finally selected. In addition, a needs assessment for early warning information was conducted through field surveys with resource managers. Based on the survey results, we decided to use four danger rating categories and six-month lead time in developing the EWS prototype by considering ASO precipitation as a trigger for forest fire. As a result, we are able to issue an ASO precipitation forecast from April to July.

3. Results and Discussions

Only individual forecast models that show consistent selection of predictors through cross and split validation procedures with significant forecast skill score (TCC) were finally selected for the EWS. The SBC method, which is based on dynamic prediction data, shows the highest model selection and is followed by statistical downscaling methods such as MWR and CIR/ITR. In most of the months, when the selected models are based on dynamic model predictions (SBC), there is a decreasing trend in TCC values as the lead times increase. Table 2 and Figure 3 show the TCC values for each month according to changes in lead time. The TCC values were calculated using MME with the condition that forecasts are issued every month. Figure 4 shows the comparison of observed and forecasted monthly precipitation for August issued in April using all selected individual models. Equal weight average of individual forecasts were used for estimating MME and the result showed a trend that lower precipitations were overestimated and higher precipitations were underestimated. Finally, monthly precipitation forecast for dry season (August to October) over four provinces in Borneo Island showed good predictability less than four-month lead time by showing temporal correlation

coefficient (TCC) greater than 0.5 in all provinces. When we consider ASO precipitation as a trigger for forest fire, we can issue an ASO precipitation forecast from April to July because we are using 6-month lead forecast data in developing prototype EWS.

In order to translate forecasted precipitation into fire danger ratings, four categories (Extreme, High, Moderate and Low) were established based on the results from the field survey. We designed a template for delivering forecast information on both precipitation and probability of forest fire for ASO period. Figure 5 shows the forecast summary for monthly precipitation and probability of forest fire in Selatan region for August to October in 1997, which was issued in April, 1997. The graph shows the graphical information for previous and current years by providing climatology (blue), observed (red), and forecasted precipitation (black). The boxplot in the figure shows the variations of predicted values by individual models. The dots at the end of the boxplots represent outliers defined by less than $Q1 - 1.5 \times IQR$ or greater than $Q3 + 1.5 \times IQR$. Where, $Q1$, $Q3$, and IQR are 25th and 75th percentiles, and difference between $Q3$ and $Q1$, respectively. The figure shows that severe drought during August to October, 1997 was closely predicted in Selatan region. The bottom-left table shows the overall summary of one-month lead forecast skill scores based on the long-term period with monitoring data. Skill score with respect to TCC according to different lead time was presented in Table 2. The used performance measures include TCC and NRMSE, which can be used for continuous variables and Accuracy and Heidke Skill Score (HSS), which in turn can be used for category forecasts. For calculating Accuracy and HSS, we equally divided the observed monthly precipitation into four categories (25% for each). Finally, the forecast information for four regions within Borneo Island will be issued each month from April to July and the forecast summary will be posted on the APCC's web hosting server (<http://www.apcc21.org/eng/html/apn.jsp>).

Month	REGION	1 MONTH	2 MONTH	3 MONTH	4 MONTH	5 MONTH	6 MONTH
JAN	Barat	0.52	0.52	0.52	0.52	0.52	0.52
	Selatan	0.69	0.69	0.63	0.49		
	Tengah	0.42	0.42	0.42	0.42		
	Timur	0.75	0.72	0.69	0.59	0.56	0.56
FEB	Barat	0.68	0.68	0.63	0.45	0.45	
	Selatan	0.72	0.68	0.59	0.5	0.5	
	Tengah						
	Timur	0.71	0.68	0.62	0.63	0.62	0.63
MAR	Barat	0.8	0.69	0.7	0.42		
	Selatan	0.82	0.82	0.76	0.55	0.55	0.45
	Tengah	0.57	0.43				
	Timur	0.68	0.69	0.67	0.59	0.57	0.55

Month	REGION	1 MONTH	2 MONTH	3 MONTH	4 MONTH	5 MONTH	6 MONTH
APR	Barat	0.68	0.68	0.63	0.43	0.43	0.43
	Selatan	0.49	0.49				
	Tengah	0.65	0.51				
	Timur	0.74	0.72	0.72	0.69	0.69	0.62
MAY	Barat	0.77	0.75	0.52	0.52	0.52	0.52
	Selatan	0.68	0.64	0.61	0.51	0.52	0.52
	Tengah	0.8	0.85	0.83	0.83	0.71	0.71
	Timur	0.65	0.67	0.72	0.72	0.57	
JUN	Barat	0.51	0.51				
	Selatan	0.58	0.58	0.41			
	Tengah						
	Timur	0.81	0.81	0.69	0.52	0.52	
JUL	Barat	0.62	0.62	0.64	0.6	0.45	0.48
	Selatan	0.6	0.59	0.58	0.58	0.47	0.47
	Tengah	0.55	0.52	0.55	0.58	0.43	0.43
	Timur	0.53	0.53	0.54	0.56	0.52	
AUG	Barat	0.7	0.69	0.68	0.67	0.67	0.65
	Selatan	0.76	0.76	0.75	0.75	0.74	0.7
	Tengah	0.67	0.66	0.67	0.68	0.68	0.62
	Timur	0.57	0.56	0.55	0.54	0.49	
SEP	Barat	0.63	0.59	0.59	0.59	0.57	0.53
	Selatan	0.54	0.54	0.53	0.51	0.52	0.54
	Tengah	0.6	0.61	0.62	0.63	0.64	0.61
	Timur	0.66	0.64	0.63	0.61	0.58	0.57
OCT	Barat	0.7	0.61	0.64	0.47	0.43	
	Selatan	0.62	0.59	0.55	0.51	0.51	0.5
	Tengah	0.6	0.57	0.53	0.53	0.46	
	Timur	0.58	0.57	0.51	0.45		
NOV	Barat	0.75	0.71	0.65	0.67	0.69	0.67
	Selatan	0.58	0.56	0.56	0.59	0.58	0.58
	Tengah	0.71	0.7	0.68	0.68	0.68	
	Timur	0.68	0.5	0.5	0.5		
DEC	Barat	0.63	0.63	0.63	0.63	0.55	0.55
	Selatan	0.69	0.62	0.62	0.62	0.62	
	Tengah	0.69	0.63	0.54	0.49	0.45	
	Timur	0.63	0.63				

TABLE 1. Temporal correlation coefficients (TCC) according to changes in lead time for predicting precipitation using multi-model ensemble (MME) average.




FIGURE 3. Temporal correlation coefficients (TCC) according to changes in lead time for predicting precipitation in Barat (a), Selatan (b), Tengah (c), and Timur (d) regions using multi-model ensemble (MME) average.





FIGURE 4. Timeseries (left) and scatter plot (right, next page) of monthly precipitation for August issued in April, where B_, M_, C_, I_ indicate SBC, MWR, CIR, ITR downscaling methods, respectively.

6-month precipitation Forecast for MAY. 1997 - OCT. 1997 (Issued: 1997-04)




Monthly skill score for JAN - DEC (1983-2007, 1-month lead forecast)

	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
TCC	0.69	0.73	0.82	0.49	0.7	0.6	0.59	0.76	0.53	0.63	0.58	0.69
NRMSE	0.75	0.8	0.74	0.86	0.76	0.8	0.79	0.66	0.86	0.76	0.8	0.71
Accuracy	0.68	0.48	0.56	0.33	0.76	0.84	0.72	0.76	0.64	0.64	0.44	0.72
HSS	0.35	0.26	0.35	-0.085	0.53	0.44	0.17	0.54	0.26	0.2	0.079	0.52

Probability of Forest Fire for 1997: Aug-Oct (%)

Extreme	5
High	95
Moderate	0
Low	0

FIGURE 5. Forecast summary for monthly precipitation and probability of forest fire in Selatan region for August to October, in 1997 (issued in April, 1997).



4. Conclusions

Based on the downscaling experiments, four different downscaling methods, in accordance with the degree of utilising the seasonal climate prediction information, were developed and integrated into the prototype of EWS in order to improve predictability. The downscaling system is based on an open source license for further training workshop and free distribution of the developed prototype. Long-term predictability of monthly precipitation for the four regions within Borneo Island was evaluated. Based on an earlier version of the prototype, APCC led a two-day workshop in Petaling Jaya, Malaysia, including hands on training sessions on statistical downscaling and prototype. Needs assessment for early warning information was also conducted through field surveys with resource managers. Finally, predictor selection algorithm in EWS prototype was improved based on the training workshop and six-month lead forecast for three months (August to October) precipitation was decided as a trigger for forest fire based on the field survey results. The SBC method, which is based on dynamic prediction data, shows the highest model selection result. In most of the months, when the selected models are based on SBC method, there is a decreasing trend in TCC values as the lead times increase. Equal weight

averages of individual forecasts were used for estimating MME and the result showed a decrease trend in yearly variation. A template was designed for delivering forecast information on both precipitation and probability of forest fire for ASO period. Monthly precipitation forecast for dry season (August to October) over four provinces in Borneo Island showed good predictability less than four-month lead time by showing TCC greater than 0.5 in all provinces. The forest fire early warning information on Southeast Asia created using the EWS will be provided through the hosting server in APCC.

Acknowledgement

We are grateful to APN for the financial support for this project and we are also grateful to the Malaysian Meteorological Department for financial and logistical support for the workshop in Petaling Jaya, Malaysia, June 9–10, 2015.

References

- Field, R. D., & Shen, S. S. P. (2008). Predictability of carbon emissions from biomass burning in Indonesia from 1997 to 2006. *Journal of Geophysical Research*, 113(G4), G04024. <http://doi.org/10.1029/2008JG000694>
- Kang, H., Park, C.-K., Hameed, S. N., & Ashok, K. (2009). Statistical downscaling of precipitation in Korea using multimodel output variables as predictors. *Monthly Weather Review*, 137(6), 1928–1938. <http://doi.org/10.1175/2008MWR2706.1>
- Kang, S., Hur, J., & Ahn, J.-B. (2014). Statistical downscaling methods based on APCC multi-model ensemble for seasonal prediction over South Korea. *International Journal of Climatology*, 34(14), 3801–3810. <http://doi.org/10.1002/joc.3952>
- Kim, M.-K., & Kim, Y.-H. (2010). Seasonal prediction of monthly precipitation in China using large-scale climate indices. *Advances in Atmospheric Sciences*, 27(1), 47–59. <http://doi.org/10.1007/s00376-009-8014-x>
- Kim, M.-K., Kim, Y.-H., & Lee, W.-S. (2007). Seasonal prediction of Korean regional climate from preceding large-scale climate indices. *International Journal of Climatology*, 27(7), 925–934. <http://doi.org/10.1002/joc.1448>